



## Effect of UV-B irradiation in *in-vitro* propagation of *Vigna mungo* cultivars

P. PERIYAKARUPPIAH<sup>1\*</sup>, M. EYINI<sup>2</sup> K. RAJARATHINAM<sup>3</sup> and M. JAYAKUMAR<sup>3</sup>

<sup>1</sup>Department of Botany, Yadava College (Autonomous), Madurai, India

<sup>2</sup>Research Centre in Botany, Thiagarajar College (Autonomous), Madurai, India

<sup>3</sup>Research Department of Botany, VHNSN College, Virudhunagar, India

\*Corresponding author E-mail: dr.p.karu@gmail.com

### Abstract

Callus induction was tried with leaf explants harvested from control and UV-B irradiated six cultivars of *Vigna mungo* (Vamban 1, Vamban 2, Vamban 3, Vamban 4, Vamban 5 and Vamban 6) to study their callus fresh weight and dry weight. Callus induction was recorded in Vamban 1, Vamban 2, Vamban 3, Vamban 4, Vamban 5 and Vamban 6 both from control and UV-B leaf explants harvested. UV-B delayed callus induction and depressed biomass accumulation. However, the callus fresh weight and dry weight were 100% compare to UV-B treated in all the cultivars of *Vigna mungo* under the control conditions. The cultivar Vamban 2 (V2) is the more sensitive one to the enhanced UV-B radiation treatment. The fresh weight and dry weight decreased by under UV-B treated plants 14%, 14%, 14%, 14%, 16% and 17% in the cultivar V1, V3, V4, V5, V6 and V2 respectively.

**Keywords:** Ultraviolet-B, *Vigna mungo*, six cultivars, Callus Initiation

### Introduction

The ozone layer in the stratosphere protects life on earth from exposure to dangerous levels of ultraviolet light. It does so by filtering out harmful ultraviolet radiation from the sun. When ozone-degrading chemicals are emitted, they mix with the atmosphere and eventually rise to the stratosphere. Chlorine and bromine catalyzes the destruction of ozone. This destruction is

occurring at a more rapid rate than ozone can be created through natural processes. The degradation of the ozone layer leads to higher levels of ultraviolet radiation reaching Earth's surface. This in turn can lead to a greater incidence of skin cancer, cataracts, and impaired immune systems, and is expected also to reduce crop yields, diminish the productivity of the oceans, and possibly to contribute to the decline of amphibious populations that is occurring around the world.

Ultraviolet-B (UV-B) radiation (280-320 nm) is a dangerous atmospheric stress (Caldwell *et al.*, 1983) as it was found to affect foliar epidermis (Bornman and Vogelmann, 1991), suppress photosynthesis (Jayakumar *et al.* 2004; Rajendiran and Ramanujam, 2004; Selvakumar *et al.* 2008; Periyakarupiah *et al.* 2015) and inhibit nodulation and nitrogen metabolism (Amudha *et al.* 2005; Vijayalakshmi and Rajendiran, 2014) in all sensitive crops. In this context, *in vitro* screening methods have to be developed to select suitable crop varieties that can survive in UV-B irradiation and also to conserve callus cells. The present study was effect of UV-B irradiation in *in-vitro* suspension culture from leaf explants of *Vigna mungo*.

## Materials and Methods

Growth of leaf calli of *V.mungo* cultivars was measured in terms of fresh and dry weight. Fresh weights of calli/explants were taken after removing the excess moisture on the surface using a blotting paper. Dry weight of callus was determined by drying the calli (1g) in a hot air oven at 60°C for 24 h. Growth was studied by determining the fresh weight and dry weight of callus after 25, 35 and 45 days of culturing. Dry weight of callus was determined after drying in a vacuum oven at 60°C until constant weight (Keskin and Kunter, 2008) for callus induction Murashige and Skoog (1962) (MS) medium supplemented with various auxins viz. Indole 3-acetic acid (IAA), indolebutyric acid (IBA), 2,4-dichlorophenoxy- acetic acid (2,4-D) and cytokinins viz. kinetin (KN), 6-benzyladenine purine (BAP) alone or in composition at different concentration (0.25, 0.5, 1, 2 mg l<sup>-1</sup>) was used for callus initiation. pH of the media was adjusted to 5.8 with 0.1 N NaOH and then agar 7.0 g l<sup>-1</sup> was added to solidify the medium and sterilized. The temperature of the growth room was maintained at 25 ±2°C. A 16 h light period/day was maintained with light intensity of 36 μmol m<sup>-2</sup> using a fluorescent light. Induced calli were transferred periodically to freshly prepared culture medium.

## Results and Discussion

Our studies showed that the greenish and yellow friable calli were observed in all the cultivars when MS medium was supplemented with 0.5 mg l<sup>-1</sup> KN and 0.5 mg l<sup>-1</sup> BAP. Enhanced UV-B exposure leads to the reduction in the fresh weight of the calli in all the cultivars for 25th, 35th and 45th days of growth. The UV-B enhanced 15 min of exposure cultivar Vamban 4, a marginal reduction of only 14% in fresh weight of callus over that of the control calli among all the six cultivars on 45th day after inoculation.

Enhanced UV-B radiation treatment decreased the calli fresh weight and dry weight in all the cultivars of *Vigna mungo* compared with control plants. Significant decrease in calli fresh and dry weight in the UV-B exposed plants calli was observed in all the six cultivars of *Vigna mungo*. Fresh weight of the calli was decreased by 14%, 14%, 14%, 14%, 16% and 17% in the cultivar V1, V3, V4, V5, V6 and V2 respectively, after 45 days of callus growth when they were irradiated with enhanced UV-B for 15 min (Table 1).

Reduction in dry weight of the calli cultivars was recorded as 6%, 6%, 7%, 8%, 8% and 19% for V3, V4, V6, V1, V5 and V2 respectively, than that of the controls after 45 days of in vitro callus growth upon receiving enhanced UV-B irradiation for 15 min. Similar reports shows that the growth rates of tea callus cultures are characteristically very low (Zagoskina *et al.*, 1990), the callus weight for both control and UV-B treatments increased significantly in the period from the 25th to 35th days of culturing. During further culturing the biomass increased only in the control callus. Hence, UV radiation facilitated earlier on completion of the growth cycle and promoted the transition of the culture to the stationary growth.

Calli grown under white light with supplementary UV radiation were also compact and dense but yellow green. The formation of chlorophyll containing cells proceeded faster in these calli than in the control culture; the difference was especially evident in the first half of the growth. In the culture irradiated with UV during growth, the accumulation pattern of phenolic compounds was clearly different.

Table 1: Influence of enhanced UV-B irradiation treatment on the Callus fresh weight and dry weight of cultivars of *V.mungo*. Values are expressed on unit fresh weight. Figures in parentheses are percentage values with reference to respective control. Mean  $\pm$  SE, n=10

Cultivar	Duration of UV-B treatment (minutes)	Callus fresh weight(F.W) and Dry weight (D.W) (g/l) Growth period (Days)					
		25		35		45	
		F.W	D.W	F.W	D.W	F.W	D.W
V1	0(Control)						
	5	51.96(100)	4.77(100)	95.07(100)	11.16(100)	127.02(100)	18.81(100)
	10	50.28(97)	4.68(98)	87.81(92)	11.1(99)	125.28(99)	18.33(97)
	15	47.49(91)	4.26(89)	81.75(86)	10.86(97)	114.6(90)	18.06(96)
V2	0(Control)						
	5	41.16(79)	4.14(87)	79.56(84)	10.74(96)	108.6(86)	17.37(92)
	10	48.83(100)	4.35(100)	91.02(100)	10.9(100)	125.32(100)	16.46(100)
	15	47.28(97)	4.23(97)	83.34(92)	10.81(99)	122.35(98)	16.34(99)
V3	0(Control)						
	5	44.47(91)	4.12(95)	78.75(87)	10.31(95)	109.54(87)	15.16(92)
	10	43.16(88)	4.11(94)	76.54(84)	10.24(94)	103.53(83)	13.36(81)
	15	49.96(100)	4.93(100)	95.75(100)	11.3(100)	127.43(100)	18.76(100)
V4	0(Control)						
	5	50.97(102)	4.82(98)	87.92(92)	11.25(100)	125.53(99)	18.52(99)
	10	47.99(96)	4.38(89)	81.83(85)	10.78(95)	114.36(90)	18.34(98)
	15	41.23(83)	4.36(88)	79.71(83)	10.75(95)	109.64(86)	17.65(94)
V5	0(Control)						
	5	52.28(100)	4.96(100)	97.07(100)	12.6(100)	131.02(100)	19.63(100)
	10	51.22(98)	4.91(99)	89.81(93)	12.53(99)	129.28(99)	19.37(99)
	15	48.41(93)	4.78(96)	83.75(86)	11.98(95)	118.65(91)	19.33(98)
V6	0(Control)						
	5	42.16(81)	4.64(94)	81.56(84)	11.94(95)	112.67(86)	18.54(94)
	10	51.34(100)	4.77(100)	95.31(100)	12.86(100)	128.02(100)	18.99(100)
	15	50.23(98)	4.53(95)	87.43(92)	11.74(91)	127.34(99)	18.67(98)
V6	0(Control)						
	5	47.27(92)	4.37(92)	81.75(86)	11.34(88)	116.65(91)	18.06(95)
	10	41.12(80)	4.34(91)	79.54(83)	11.33(88)	110.63(86)	17.56(92)
	15	51.93(100)	4.79(100)	95.56(100)	11.19(100)	127.65(100)	18.85(100)
V6	0(Control)						
	5	50.25(97)	4.72(99)	87.76(92)	11.12(99)	125.26(98)	18.34(97)
	10	47.42(91)	4.28(89)	81.87(86)	10.83(97)	114.65(90)	18.32(97)
	15	41.11(79)	4.16(87)	79.34(83)	10.43(93)	107.76(84)	17.54(93)

The largest total content of soluble phenolic compounds was noted by the end of the growth in plants (Zagoskina *et al.* 2005; Nagarajan *et al.* 2008; Benjamin *et al.* 2010; Periyakaruppiyah *et al.* 2012a; Periyakaruppiyah *et al.* 2012b; Periyakaruppiyah *et al.* 2014). Enhanced UV-B exposure leads to the reduction in the number of callus cells. The suspension cells were 100% viable in all the cultivars of *Vigna mungo* under the control conditions. The cultivar Vamban 2 (V2) is the more sensitive one to the enhanced UV-B radiation treatment. The suspension cells which

received 10 and 15 mins of enhanced UV-B radiation only 53% and 38% cells were viable, respectively (Periyakaruppiyah *et al.* 2016).

## Conclusion

The *in vitro* culture is a very attractive technique for the cultivation of callus mass, particularly fresh and dry weight of *Vigna mungo* cultivars. In conclusion, clearly UV-B irradiation could play a role in leaf explants. Marginal decrease in plant biomass under enhanced UV-B radiation conditions as observed in *Vigna mungo* cultivar Vamban 3 and 4 may be of great significance from *in vitro* propagation.

## References

Amudha, P., Jayakumar, M. and Kulandaivelu, G 2005 Impacts of ambient solar UV (280-400nm) radiation on three tropical legumes. *J. Plant Biol.* 48: 284-291

Benjamin Jeya Rathna Kumar, P. Periyakaruppiyah, P. Jeyakumar, S. Rajarathinam K and Jayakumar M. 2010 Influence of enhanced UV-B (285-320 nm) radiation on secondary metabolites in *Justicia gendarussa* Burm.f.-a medicinal plant. *Int. Conference on Medicinal Plants and Herbal Drugs: Challenges and Opportunities in cultivation Sustainable Utilization and Conservation*

Bornman JF and Vogelmann TC 1991 Effect of UV-B radiation on leaf optical properties measured with fibre optics. *J. Experimental Botany.* 42: 647-554

Caldwell M, Gold G, Harris G and Ashurst W 1983 A modulated lamp system for solar UV-B (280-320 nm) supplementation studies in the field. *Photochemistry and Photobiology.* 37: 479-485

Jayakumar, M., Amudha, P. and Kulandaivelu, G 2004 Effect of low doses of UV-A (320-400 nm) and UV-B (280-320 nm) radiation on photosynthetic activities in *Phaseolus mungo* L. *J. Plant Biol.* 47: 105-110

Keskin, N. and Kunter, B. 2008 Production of transresveratrol in Cabernet sauvignon (*Vitis vinifera* L.) callus culture in response to ultraviolet-C irradiation. *Vitis.* 47: 193-196

Murashige T and Skoog F 1962 A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiologia Plantarum*. 15: 473-497

Nagarajan, P. Periyakaruppiyah, P. Premkumar, G. Muthuramkumar, S. Rajarathinam K and Jayakumar. M 2008 Effect of enhanced UV-B (280-320nm) radiation on the changes in the secondary metabolites content in *Vitex negundo*- a medicinal plant. *National Conference on Cultivation, Processing and Bioprospecting of Ethnomedicine*

Periyakaruppiyah, P. Eyini, M. Rajarathinam K and M. Jayakumar 2016 Effect of UV-B (280-320nm) radiation in *IN-VITRO* culture from callus explant of black gram cultivars. *Int. J. Current Sci. Res.* 2(4): 518-523

Periyakaruppiyah, P. Jayakumar M and Rajarathinam K. 2012 a Effect of UV-B on enhancement of Anthocyanin and Flavonoids contents in the Different Cultivars of *Vigna mungo*. *National seminar on Herbal Wealth-Traditions and New Approaches*

Periyakaruppiyah, P. Nagarajan, P. Rajarathinam K and Jayakumar M 2012 b Effect of enhance UV-B (280-320 nm) radiation on the changes in the secondary metabolites content in *solanum trilobatum*, a medicinal plant. *J. Theoretical and Experimental Biology*. 9: 53-58

Periyakaruppiyah, P. Rajarathinam, K. Eyini M and Jayakumar M 2015 Effect of enhanced UV-B (280-320nm) radiation on the changes in the Allelochemicals in *Solanum trilobatum*- a medicinal plant *In: Current and Emerging Trends in Biological Sciences* (Eds) P.Annapoorani R.Renuka and P. Mehalingam, Publishing Virudhunagar Hindu Nadars' Senthikumara Nadar College (Autonomous), pp. 169-177

Periyakaruppiyah, P. Yasothkumar N. Eyini M and Jayakumar M. 2014 Effect of enhanced UV-B irradiation on photosynthetic pigments and non- photosynthetic pigments of different cultivars of *Vigna mungo*. *National seminar on Biovision – 2014*

Rajendiran K and Ramanujam MP 2004 Improvement of biomass partitioning, flowering and yield by triadimefon in UV-B stressed *Vigna radiata* (L.) Wilczek. *Biologia Plantarum* 48: 145-148

Selvakumar, V. Periyakaruppiyah, P. Rajarathinam K and Jayakumar M. 2008 Effect of UV-B (280-320 nm) on the Enhancement of Allelochemicals (Phenols) in Catharanthus roseus Plants. *5th World Congress on Allelopathy*. Abstract No.160

Vijayalakshmi R and Rajendiran K 2014 Impact of ultraviolet-B radiation on nodulation and nitrogen metabolism in Phaseolus vulgaris L. cv. Prevail. *Int. J. Advanced Biological Res.* 4(3): 339-342

Zagoskina, N.V., Alyavina, A.K., Gladyshko, T.O., Lapshin, P.V., Egorova, E.A. and Bukhov, N.G. 2005 Ultraviolet rays promote development of photosystem II photochemical activity and accumulation of phenolic compounds in the tea callus culture (*Camellia sinensis*). *Rus. J. Plant Physiol.*, 52: 731-739

IJCSR Specialities

**\$ Impact Factor – IBI – 2.9; GIF – 0.676 & SIF – 0.54**

**\$ Indexed over 30 databases**

**\$ Monthly Issue**

<http://www.drbgpublications.in/ijcsr.php>