



Growth Judgment of *Channa striatus* fry fed with different live feeds

D.Kumar^{1*}, M.A.Haniffa²

¹Department of Zoology, V.H.N.S.N. College (Autonomous), Virudhunagar, India

²Centre for Aquaculture Research and Extension (CARE), St. Xavier's College, Palayamkottai, Tirunelveli, India

*Corresponding author email: drkumarbio@yahoo.com

Abstract

Today live feed play a very specific role in the aquaculture industry. Various live feed organisms were distributed throughout the world. The live feed which enhance the growth of the fish larval stage and as well as to maintain the survival of the fish population. In the initial stage of the exogenous feeding the young ones, which advice to provide plankton soup in a very small size its depending on the mouth size of the young ones. In our study, I have chosen the Murrels species *Channa striatus* which is commercially important fish one in India, That's why we prepared that species and provided the locally available live feed viz; Tubifix (D1) Chironomus larvae (D2) Beef liver (D3) Mosquito larvae (D4) and Plankton (D5) fed to the *Channa striatus* fry over a period of 45 days. Among the food *Chironomus larvae* was found to produce the best SGR (1.529 ± 0.052) Weight gain (919.00 ± 62.324) RGR % (9.162 ± 0.70) ADG (g/day) (0.0874 ± 0.0013) Growth rate (%) (40.55 ± 0.299) Survival % (96.66 ± 4.714) and the poor growth parameters were recorded in Plankton soup (D5) respectively.

Keywords: *Channa striatus*, growth, nutrition, live feeds.

Introduction

Channa striatus snakehead is an air breathing, and carnivorous in nature. They are widely distributed in Africa and Asia. It is having high market due to its tasty and flesh. They support economically important fisheries and aquaculture industry in many Asian countries

(Ling 1979, Chen 1990). Among murrels *C.striatus* forms a significant role in capture fisheries of India. Characteristics of this fish that make it desirable cultivable food fish include rapid growth and the ability of the fish to store and use atmospheric oxygen for respiration in waters with low dissolved oxygen and they can withstand higher stocking densities also. It has been estimated that out of 18,000 t of marketable surplus air breathing fishes caught from natural resources in India, murrels account for nearly 12,000 t (Jhingran 1975) with major part of them constituted by *Channa marulius*, *Channa striatus* and *Channa punctatus*. However, murrel culture is not practiced in a well-defined way in India due to several reasons. One of these is that there are no seed supply / sales center for murrels in country.

Various dry feed formulae have been investigated as possible substitutes of live food for larval development (Appelbaum and Dor, 1978; Dabrowski 1983; 1984). In recent years suitability of various dry feed formulae has been investigated for the rearing of Cyprinid and catfish larvae (Bryant and Matty, 1981; Msiska, 1981; Hecht and Viljoen, 1982). However it has been shown that formulated compound diets do not provide optimal larval growth when used exclusively as larval food, especially during the early larvae stages of Cyprinids and catfish (Dabrowski 1984; Prinsloo and Schoonbee, 1986), therefore live food provide a substantial availability of protein and other essential nutrients (Jhingran, 1975; Ahmed, 1994, Thakur, 1978, Munnet 1979) provided some basic information on the feeding of *C.batrachus* fry. However development of suitable feed for rearing *Channa striatus* fry is lacking. It is therefore important to study the efficacy of a few selective feeds.

The fish farmers therefore depend on wild collection, which are unpredictable. Further, the rearing of hatchlings, post larvae and fry of *C.striatus* is a complicated process unlike the raising of carp fry, which has been standardized in some extent. Recently attempts have been made on larval nutrition of *Channa striatus* by (Qin *et al.*, (1997), Samantaray and Mohanty 1997). But these authors have provided formulated pelleted feeds (instead of live feed) to the larvae resulting in poor survival and growth. In the present study a comparison is made on the growth of *C.striatus* larvae using tubifix, Chironomus, plankton and Beef liver and mosquito larvae as alternative live foods during the early larval growth phase of this species.

Materials and Method

Channa striatus fry (length: 3.57±0.05cm, weight, 0.425±0.03 g) were collected from (CARE) earthen pond and acclimatized in the cement tanks for a period of one week during this period they were fed with plankton soup. They were grouped into 5 batches and stocked for 10 individual for each treatment with three replicates and reared in plastic troughs. (Capacity 15/Lt).

Water quality parameters viz; temperature 29°C ±1°C dissolved oxygen 6.1-6.6 mg/lit and pH 7.5-8.25 were recorded throughout the study. They were fed with Tubifix, Chironomus larvae, Beef liver Mosquito larvae, and Plankton soup twice/day (11.00hrs, 13.00hrs) *ad libitum*. The feeding trial was continued for a period of 45 days. Water was changed every day with minimal disturbance to the experimental animals full nos. the length, and weight were recorded once in every fortnight, the growth parameters viz, weight gain (%), specific growth rate and (%/day) and survival (%) were estimated.

Formula

The energy budget of the fish was calculated by following the IBP formula

$$\text{Specific growth rate SGR } (\%/day) = \frac{\ln \log \text{ Final body weight} - \ln \log \text{ initial live weight (g)}}{\text{Experimental duration}} \times 100$$

$$\text{Weight gain } (\%) = \frac{\text{Final live weight} - \text{Initial live weight}}{\text{Days}} \times 100$$

$$\text{Average daily growth rate ADG } (\%) = \frac{\text{Growth (live wt. g)}}{\text{Duration}} \times 100$$

$$\text{Survival } (\%) = \frac{\text{No of fish introduced}}{\text{No of fish survived}} \times 100$$

$$\text{Mean growth rate} = \frac{\text{Final mean weight}}{\text{Initial mean weight/days}}$$

Results

All foods were readily accepted from the start of feeding. The beef liver was taken by larvae from the bottom of the trough, Where as Zooplankton, mosquito larvae in the mid water of the trough and the tubifix, Chironomus taken by the larvae from the bottom. The growth parameter and body composition of the fish fed on different diets and was presented in table 1 and from the table 2 body proteins, carbohydrate and lipid of the fish is higher in the diet of D2 and followed by other diets groups. And fig 1 shows the Specific growth rate and relative growth rate and growth rate of *Channa striatus* fry

Live food is regarded as the best for fishes, many crustacean, insect larvae. In present study the result revealed that the highest SGR (1.529 %/day), was obtained in the individual fed with Chironomus larvae (Table 1) followed by tubifix fed on (1.383%/day), beef liver (1.22%/day) mosquito larvae 1.112%/day and plankton (1.082%/day). The best weight gain was obtained those fed on Chironomus larvae (919.00). And followed by (705.64) those fed on tubifix. The best survival rate of (96.66 %) was obtained in *Channa striatus* fed on Chironomus larvae whereas the lowest survival (73.33 %) were observed in those fed on plankton.

Discussion

The results showed that live food, in particular Chironomus larvae is a most desirable diet for the rearing of the *Channa striatus* larvae. The importance of artemia as live food (Msiska, 1981) is again confined by this investigation, However Prinsloo and Schoonbee, (1986) observed zooplankton as best live food in comparison to commercial dry feed for the rearing of the silver carp and grass carp species over a period of 10-14 days; silver carp and grass carp larvae accured relatively better growth with zooplankton as compared to commercial dry food. In our present study instead of Zooplankton Chironomus displayed superior growth and other parameter were recorded, due to the higher protein 61.17% and haem content of the Chironomus larvae. Qin *et al.*, (1997), also developed a protocol for weaning larval snake heads from live artemia to formulated feed, but grow out performance with formulated feed was not evaluated.

Live food is an important diet in the rearing of larvae of a member of fish species (Msiska, 1981; Stenson, 1982) indicated the importance of rotifer *Brachionus plicatilis* for mass larval rearing of fishes and stressed the value of the inclusion of rotifers in combination with

artificial dry feeds for the optimum growth of *Cyprinus carpio* larvae. Matlak and Matlak (1976) indicated that rotifers are important food items of carp larvae during the first three weeks in nursery ponds. Zooplankton is the best larval food for a variety of fish larvae (Kilambi and Zdinak, 1982; Geiger, 1983a, and Dabrowski, 1984). The nutritional value of artemia for *Cyprinus carpio* larvae indicated good growth (Bryant and matty 1981). A variety of dry foods were used for the rearing of *C. carpio* larvae (Appelbaum and Dor 1978; Hecht and Viljean, 1982).

According to (Cahu *et al.*, (1998), larvae receiving live food also showed better survival and growth than larvae receiving artificial diets, sea bass *Dicentrarchus labrax*, Abi-Ayad and Kestemont, (1994) observed the highest SGR in gold fish *Carassius auratus* larvae fed the mixed diets. Whereas the lowest SGR (1.325%/day) fed on plankton. Numbers of reports are available on larval rearing, Mahseer larvae, (Rai, 1990), also reported that larvae of Mahseer fed with plankton showed better growth when fed with other supplementary diets. In addition to the fry of katle *Acrossocheilus hexagonolepis* also showed better growth, when fed with 30% protein content with plankton soup (Rai, 1990).

Malkotra and Munchi, (1985) found that formulated feed might also be physically unsuitable for most first feeding fish larvae because large food particles that didn't pass down the gut could subject larvae to physical stress or physiological stress. The fish growth rate is generally related to availability and density of optimal food (Mittelbach, 1981). Walleye growth increased after they switched their diets from zooplankton to Chironomids (Fox *et al.*, 1989). But in our present study Chironomus larvae displayed superior growth fed with the *Channa striatus* fry is reported.

Dabrowski, (1982) reported that many small fish larvae do not have the enzymes for digesting non-living diets. So we were applied different types of feeds. *Artemia nauplia* suitable for rearing of the young ones. Some relevant reports are available; Fluechter (1980) found that protein digestion enzymes in live *Artemia nauplia* were responsible for successful rearing of White fish (*Coregonus lavaretus*) larvae.

Conclusion

Our present results strongly support the use of live food in the early larval growth phase of *C. striatus* fry. So the blood worm is strongly recommended to the rearing of *C. striatus* larvae as well as fry.

Reference

- Abi-Ayad, A. and P. Kestemont 1994. Comparison of the nutritional Status of gold fish *Carassius auratus* larvae with live mixed or dry diet. *Aquaculture*. 128: 163-176
- Ahmed, G.U. 1994. Effect of first food on the growth of African catfish (*Clarias gariepinus*) during the primary nursing phase. *BAV. Res. Prog.* 8: 567-574
- Appelbaum, S. and U. Dor 1978. Zehntagige Anfütterung von Karpfenlarven (*Cyprinus carpio*) mit Trockenfutter auf Kommerzieller basis. *Der Fischwirt*. 28(5): 25-26
- Bryant, P.L. and A.J. Matty 1981. Adaptation of carp (*Cyprinus carpio*) larvae to artificial diets 1: Optimum feeding rate and adaptation age for a commercial diet. *Aquaculture*. 23: 275-286
- Cahu, C, J.h Zambonino infante A.M. Escatfre. P. Bergot and S. Kaushik 1998. Preliminary results on seabass (*Dicentrarchus labrax*) larvae rearing with compound diet from first feeding. Comparison with carp (*Cyprinus carpio*) larvae. *Aquaculture*. 169: 1-7
- Chen, L.C. 1990. Snake head culture: 39-42. In L.C. Chen (ed). *Aquaculture in Taiwan*. black well scientific publication, London
- Dabrowski, K. 1982. Proteolytic enzyme activity decline in starving fish alevins and larvae *Environ. Biol. Fish.* 7: 73-76
- Dabrowski, K. 1983. Comparative aspect of protein digestion and amino acid absorption in fish and other animals. *Com. Biochem. Physiol.* 74 A: 417-425
- Dabrowski, K. 1984. Influence of initial weight during the change from live to compound feed on the survival and growth of four cyprinids. *Aquaculture*. 148: 105-113
- Fluechter, J. 1980. Revive of the present knowledge of rearing white fish (Coregonidae) larvae. *Aquaculture*. 19: 191-208
- Fox M.G Keast J.a swainson, R.J. 1989. The effect of fertilization regime on juvenile Walleye growth and Prey utilization in rearing Ponds. *Environ. Bio. Fish.* 26: 129-142
- Geiger. J.G. 1983a. Zooplankton production and manipulation in striped bass rearing pond. *Aquaculture*. 35: 331-351

- Hecht, Y. and H. Viljoen 1982. Observation on the suitability of various dry feeds for the commercial rearing of carp, *Cyprinus carpio* larvae in South Africa. Water SA. 8(1): 58-65
- Hoogendoorn, H. 1980. Controlled propagation of the African catfish, *Clariass lazera* (C & V). III: Feeding and growth of fry. Aquaculture. 21: 233-241
- Jhingran 1975. Fish and fisheries of India. Hindustan publishing corporation 954 pp.
- Kilambi, R.V. and A. Zdinak. 1982. Food intake and growth of hybrid carp (female carp, *Ctenopharyngodon idella* and male bighead, *Aristichthys (Hypothalmichthys nobilis)* fed on zooplankton and chara. J. Fish Biol: 21: 63-67
- Ling, S.W. 1979. Snake head *Ophiocephalus spp.* Culture, pp.60-64 In, S.W.Ling (ed) Aquaculture in south east Asia. A historical review. University of Washington press, Seattle
- Malkotra, Y.R. munshi. S; 1985. first feeding and survival of Aspido Paria Morar (Cyprinidae) Trans.AM.Fish. Soc.114: 286-290
- Matlak, J. and O. Matlaak 1976. The natural food of carp fy (*Cyprinus carpio*). Acta.Hydrobiol. 18(3): 203-228
- Mittelbach G.G 1981. Foraging efficiency and body size; a study of optimal diet and habitat use by blue gills, Ecology. 62: 1370-1386
- Msiska, O.V. 1981. Rearing of the fry of the African catfish *Clarius lazera* (C & V) using live and artificial feed stuffs. Bamidgeh: 33: 122-127
- Munnet, S.K. 1979. An experiment of feeding of magur, *Clarius batrachus* (linn). J. Inland fish. Soc. India 11(2): 10-14
- Prinsloo, J.F. and J.H. Schoonbee 1986. Comparison of the early larval growth rates of the Chinese silver carp *Hypophthalmichthys nobilis* using live and artificial feed. Water SA: 12(4) 229-234
- Qin, J., Fast, A.N., De Anda D. and Weidenbach R.P 1997. Growth and survival of larval snake head *Channa striatus*, fed different diets. Aquaculture. 148: 105-113
- Rai, A.K. 1990. Propagation of katle (*Acrossocheilus hexagonolepis*) and studies on food habits and growth of young ones in different dietary protein level. A project report submitted

to International development research Centre (IDRC) lalitpur, nepal inland fisheries Dev.Project, Harihar Bhawan, Pulchowk pp 22

Samantaray. K and S.S Mohanty 1997. Interactions of dietary levels of protein and energy on fingerling snakehead, *Channa striatus*. Aquaculture. 156 pp 241-249

Stenson, J.A.E. 1982. Fish impact on rotifer community structure. Hydrobiologia. 87: 57-64

Thakur, N.K., 1978. On the food of the air breathing catfish *Clarias batrachus* (Linn). occurring in wild water. Hydrobiologia. 63(3): 421-431

Table 1: Growth and Survival of *Channa striatus* fry fed on different types of diets

Parameters	Diets				
	Tubifix (D1)	Chironomus larvae (D2)	Beef liver (D3)	Mosquito (D4)	Plankton (D5)
Initial length (cm)	3.506±0.024	3.603±0.018	3.586 ±0.109	3.66±0.141	3.523±0.053
Initial weight (mg/g)	0.406±0.020	0.390±0.031	0.425±0.041	0.443±0.040	0.463±0.026
Final length (cm)	6.918±0.072	7.66±0.082	6.69±0.052	6.512±0.305	6.054±0.069
Final weight (mg/g)	3.267±0.020	3.948±0.056	2.85±0.081	2.11±0.082	2.126±0.010
Experimental duration	45	45	45	45	45
SGR %/day	1.383±0.049	1.529±0.052	1.22±0.097	1.112±0.081	1.082±0.045
Weight gain %	705.64±41.805	919.00±62.324	578.15±74.78	380.20±38.37	360.185±26.47
ADG (g/day)	0.0723±0.0004	0.0874±0.0013	0.0629±0.0016	0.0466±0.001	0.0472±0.0002
Growth rate (%)	39.39±0.301	40.55±0.299	37.79±0.600	35.56±0.760	35.18±0.551
Survival %	90±14.14	96.66±4.714	76.66±4.714	66.66±4.714	73.33±9.428

Table 2: Body composition of the *Channa striatus* fry fed on different diets

Parameters	Diets					
	Initial	D1	D2	D3	D4	D5
Protein %	14.01	15.14	16.28	15.68	14.21	14.32
Carbohydrate %	0.88	0.98	1.01	0.88	0.94	0.93
Lipid %	3.14	3.16	3.06	2.94	3.01	2.84
Ash %	3.2	3.5	3.9	2.8	3.2	3.4
Moisture %	76.14	74.15	73.48	75.14	74.28	75.01

D1-Tubifix, D2-Chironomus, D3- Beef liver, D4-Mosquito, D5- Plankton.

IJCSR specialities

\$ **Systematic Impact Factor (SIF) – 2.306**

\$ PubMed: NLM id: 101671355

\$ **THOMSON REUTERS – RESEARCHERID - [M-7259-2015](#)**

\$ **IJCSR Journal ISRA: RUN Value: 30.10.2015.797**

\$ **Journal – DOI: 05.2016-31641248**

\$ Search engines have verified that **14,500 Cities** in **109 countries**

\$ *Monthly Issue*

<http://www.drbgpublications.in/ijcsr.php>